

Massachusetts Institute of Technology  
Physics Department

Physics 8.20  
Special Relativity  
Room 2-190

IAP 2003  
January 21, 2003  
7:30–9:30 pm

## Midterm

Your name: \_\_\_\_\_

### Instructions

- You have two hours for this test. Papers will be picked up at 9:30 pm sharp.
- The exam is scored on a basis of 100 points.
- Please do ALL your work on the pages of the exam. There are extra pages stapled at the end if you run out of space.
- Please remember to put your name on each page of the exam.
- You may use the single page of notes that you have prepared.
- If you have any questions please feel free to ask the proctor(s).

**Information**

$c \approx 300\text{km/millisecond}$

Lorentz transformation (along the  $x$ -axis) and its inverse

$$\begin{array}{ll} x' = \gamma(x - \beta ct) & x = \gamma(x' + \beta ct') \\ y' = y & y = y' \\ z' = z & z = z' \\ ct' = \gamma(ct - \beta x) & ct = \gamma(ct' + \beta x') \end{array}$$

where  $\beta = v/c$ , and  $\gamma = 1/\sqrt{1 - \beta^2}$ .

Velocity addition (relative motion along the  $x$ -axis):

$$\begin{aligned} u'_x &= \frac{u_x - v}{1 - u_x v/c^2} \\ u'_y &= \frac{u_y}{\gamma(1 - u_x v/c^2)} \\ u'_z &= \frac{u_z}{\gamma(1 - u_x v/c^2)} \end{aligned}$$

Doppler shift

Longitudinal

$$\nu = \sqrt{\frac{1 + \beta}{1 - \beta}} \nu_0$$

Quadratic equation:

$$\begin{aligned} ax^2 + bx + c &= 0 \\ x &= \frac{1}{2a} \left( -b \pm \sqrt{b^2 - 4ac} \right) \end{aligned}$$

Binomial expansion:

$$(1 + a)^b = 1 + ba + \frac{b(b-1)}{2} a^2 + \dots$$

1. Problem 1 [27 points] Short Answer Questions

*The equations below should be answered briefly, no more than five lines and no calculations. Please be brief and to-the-point.*

- (a) [3 points] Define an inertial frame and give an example of a *non*-inertial frame.
- (b) [3 points] Critique the statement: “If two events are simultaneous in one inertial frame then they are simultaneous in all inertial frames.”
- (c) [3 points] Relative to any event, spacetime can be decomposed into *future*, *past*, and *elsewhere*. What, precisely, is the significance of the region called “elsewhere”?

Your Name: \_\_\_\_\_

Physics 8.20 Midterm, IAP 2003 4

(d) [3 points] In pre-Einstein physics the concept of an “incompressible fluid” refers to a fluid which cannot be compressed by any force at hand. Is this concept consistent with special relativity? Explain.

(e) [3 points] Light is emitted from a star that moves perpendicular to your line of sight at a velocity close to  $c$ . Is it redshifted or blueshifted or neither?

(f) [3 points] Define proper time.

[Continued on following page]

**Your Name:** \_\_\_\_\_

*Physics 8.20 Midterm, IAP 2003* 5

(g) [3 points] In one sentence summarize the fundamental result of the Michelson-Morley experiment.

(h) [3 points] An intense searchlight on the ground sweeps across the sky. A layer of clouds overhead reflects the light from the searchlight beam back to an observer standing next to the search light. Evaluate the following statement: “The spot made by the searchlight as seen by the observer can move faster than the speed of light”

**2. Problem 2 [15 points] The Concept of Relativity**

The concept of relativity:

*The laws of Mechanics are such that they hold in all inertial frames*

was known in mechanics before Einstein, and was based on the “Galilean” transformation,

$$\vec{x}' = \vec{x} - \vec{v}t, \quad t' = t$$

relating the coordinates of inertial observers moving at a relative velocity  $\vec{v}$ . Which of the following force laws listed below are consistent with this pre-Einstein concept of relativity? Give a *short* explanation.

[Labels 1 and 2 refer to two different bodies.  $\vec{x}$ ,  $\vec{v}$ , and  $\vec{a}$  are the position, velocity, and acceleration, respectively.  $k$ ,  $b$ , and  $g$  are positive constants.]

(a)  $m_1\vec{a}_1 = -k\vec{x}_1$

(b)  $m_1\vec{a}_1 = -b\vec{v}_1$

(c)  $m_1\vec{a}_1 = -k(\vec{x}_1 - \vec{x}_2)$

**3. Problem 3 [18 points] Relations between two events**

Two events,  $A$  and  $B$ , occur on the  $x$ -axis and *at the same time* in the frame  $\Sigma$ . In that frame their separation is  $\Delta x_0$

- (a) [4 points] Show that the two events occur at *greater* separation in any other inertial frame,  $\Sigma'$ , moving in the  $\hat{x}$  direction relative to  $\Sigma$ .
- (b) [4 points] Although  $A$  and  $B$  are simultaneous in  $\Sigma$ , they occur at different times in frames moving with respect to  $\Sigma$ . Show that it is possible to find a frame  $\Sigma'$ , (moving along the  $\hat{x}$ -direction relative to  $\Sigma$ ) in which event  $A$  occurs *arbitrarily long before* event  $B$ .
- (c) [10 points] Define the “invariant interval”,  $\Delta S^2$ , and use the Lorentz transformation from  $\Sigma$  to  $\Sigma'$  to prove that it is indeed invariant, ie. prove  $\Delta S^2 = \Delta S'^2$ . Use the invariant interval to show that in a frame where event  $A$  occurs very long before event  $B$ , it also occurs very far away from event  $B$ .

Your Name: \_\_\_\_\_

Physics 8.20 Midterm, IAP 2003 8

**4. Problem 4 [20 points] Particle decay**

A  $\rho$ -meson decays into two pions. In the rest frame of the  $\rho$  the pions are emitted (back-to-back), each with speed  $0.7c$ .

- (a) [6 points] What is the speed of one pion relative to the other?
- (b) [7 points] Suppose the decay is observed in a frame, which we'll call the "lab frame", where the initial  $\rho$  has speed  $0.9c$  before it decays. In the lab frame, what is the greatest speed that one of the pions can have?
- (c) [7 points] Same conditions as (b). What is the least speed a pion can have?

**5. Problem 4 [20 points] Trip to a nearby planet**

Suppose a very interesting planet has been discovered near a star 50 light years from the Earth. In the year 2050 NASA decides to send an expedition to explore this planet. The decision is made that the men and women on the expedition should reach the planet after 20 years of their own body time has elapsed.

- (a) [2 points] What should be the speed at which the expedition's ship travels from Earth to the planet? [You may assume constant speed and ignore initial acceleration and deceleration.]
- (b) [3 points] How long does the expedition appear to take according to the NASA observers on Earth?
- (c) [5 points] The ship carries radio transmitters tuned to the frequency 90 megahertz ( $9 \times 10^7$  cycles per second) that they use to send messages back to earth. What frequency should the NASA base station tune to in order to receive these signals?
- (d) [10 points] When the ship reaches the half way point, the astronauts get impatient and send out an advanced party. The advanced party leaves the ship with a relative velocity of  $0.99c$ . [Again ignore acceleration and deceleration.] How many years have passed since the original departure from Earth *according to the astronauts in the advanced party* when they reach the planet?