

## Exercises on differential equations and $e^{At}$

**Problem 23.1:** (6.3 #14.a *Introduction to Linear Algebra*: Strang) The matrix in this question is skew-symmetric ( $A^T = -A$ ):

$$\frac{d\mathbf{u}}{dt} = \begin{bmatrix} 0 & c & -b \\ -c & 0 & a \\ b & -a & 0 \end{bmatrix} \mathbf{u} \quad \text{or} \quad \begin{aligned} u_1' &= cu_2 - bu_3 \\ u_2' &= au_3 - cu_1 \\ u_3' &= bu_1 - au_2. \end{aligned}$$

Find the derivative of  $\|\mathbf{u}(t)\|^2$  using the definition:

$$\|\mathbf{u}(t)\|^2 = u_1^2 + u_2^2 + u_3^2.$$

What does this tell you about the rate of change of the length of  $\mathbf{u}$ ? What does this tell you about the range of values of  $\mathbf{u}(t)$ ?

**Solution:**

$$\begin{aligned} \frac{d\|\mathbf{u}(t)\|^2}{dt} &= \frac{d(u_1^2 + u_2^2 + u_3^2)}{dt} \\ &= 2u_1u_1' + 2u_2u_2' + 2u_3u_3' \\ &= 2u_1(cu_2 - bu_3) + 2u_2(au_3 - cu_1) + 2u_3(bu_1 - au_2) \\ &= 0. \end{aligned}$$

This means  $\|\mathbf{u}(t)\|^2$  stays equal to  $\|\mathbf{u}(0)\|^2$ . Because  $\mathbf{u}(t)$  never changes length, it is always on the circumference of a circle of radius  $\|\mathbf{u}(0)\|$ .

**Problem 23.2:** (6.3 #24.) Write  $A = \begin{bmatrix} 1 & 1 \\ 0 & 3 \end{bmatrix}$  as  $S\Lambda S^{-1}$ . Multiply  $Se^{\Lambda t}S^{-1}$  to find the matrix exponential  $e^{At}$ . Check your work by evaluating  $e^{At}$  and the derivative of  $e^{At}$  when  $t = 0$ .

**Solution:** The eigenvalues of  $A$  are  $\lambda_1 = 1$  and  $\lambda_2 = 3$ , with corresponding eigenvectors  $\mathbf{x}_1 = (1, 0)$  and  $\mathbf{x}_2 = (1, 2)$ . This gives us the following values for  $S, \Lambda$ , and  $S^{-1}$ :

$$S = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}, \Lambda = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}, S^{-1} = \begin{bmatrix} 1 & -1/2 \\ 0 & 1/2 \end{bmatrix}.$$

We use these to find  $e^{At}$  :

$$Se^{At}S^{-1} = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} e^t & 0 \\ 0 & e^{3t} \end{bmatrix} \begin{bmatrix} 1 & -1/2 \\ 0 & 1/2 \end{bmatrix} = \begin{bmatrix} e^t & .5e^{3t} - .5e^t \\ 0 & e^{3t} \end{bmatrix} = e^{At}.$$

Check:

$$e^{At} = \begin{bmatrix} e^t & .5e^{3t} - .5e^t \\ 0 & e^{3t} \end{bmatrix} \text{ equals } I \text{ when } t = 0. \checkmark$$

$$\frac{de^{At}}{dt} = \begin{bmatrix} e^t & 1.5e^{3t} - .5e^t \\ 0 & 3e^{3t} \end{bmatrix}.$$

$$\left. \frac{de^{At}}{dt} \right|_{t=0} = \begin{bmatrix} 1 & 1 \\ 0 & 3 \end{bmatrix} = A. \checkmark$$