

Exercises on Markov matrices; Fourier series

Problem 24.1: (6.4 #7. *Introduction to Linear Algebra*: Strang)

- a) Find a symmetric matrix $\begin{bmatrix} 1 & b \\ b & 1 \end{bmatrix}$ that has a negative eigenvalue.
- b) How do you know it must have a negative pivot?
- c) How do you know it can't have two negative eigenvalues?

Solution:

- a) The eigenvalues of that matrix are $1 \pm b$. If $b > 1$ or $b < -1$ the matrix has a negative eigenvalue.
- b) The pivots have the same signs as the eigenvalues. If the matrix has a negative eigenvalue, then it must have a negative pivot.
- c) To obtain one negative eigenvalue, we choose either $b > 1$ or $b < -1$ (as stated in part (a)). If we choose $b > 1$, then $\lambda_1 = 1 + b$ will be positive while $\lambda_2 = 1 - b$ will be negative. Alternatively, if we choose $b < -1$, then $\lambda_1 = 1 + b$ will be negative while $\lambda_2 = 1 - b$ will be positive. Therefore this matrix cannot have two negative eigenvalues.

Problem 24.2: (6.4 #23.) Which of these classes of matrices do A and B belong to: invertible, orthogonal, projection, permutation, diagonalizable, Markov?

$$A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \quad B = \frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

Which of these factorizations are possible for A and B : LU , QR , $S\Lambda S^{-1}$, or $Q\Lambda Q^T$?

Solution:

a) For A :

$$\det A = -1 \neq 0.$$

A is **invertible**.

$$AA^T = I.$$

A is **orthogonal**.

$$A^2 = I \neq A.$$

A is **not a projection**.

A has one 1 in each row and column with 0's elsewhere.

A is a **permutation**.

$$A = A^T, \text{ so } A \text{ is symmetric.}$$

A is **diagonalizable**.

Each column of A sums to one.

A is **Markov**.

All of the factorizations are possible for A : LU and QR are always possible, $S\Lambda S^{-1}$ is possible because it is diagonalizable, and $Q\Lambda Q^T$ is possible because it is symmetric.

b) For B :

$$\det B = 0.$$

B is **not invertible**.

$$BB^T \neq I.$$

B is **not orthogonal**.

$$B^2 = B.$$

B is a **projection**.

B does not have one 1 in each row and each column, with 0's elsewhere.

B is **not a permutation**.

$$B = B^T \text{ so } B \text{ is symmetric.}$$

B is **diagonalizable**.

Each column of B sums to one.

B is **Markov**.

All of the factorizations are possible for B : LU and QR are always possible, $S\Lambda S^{-1}$ is possible because it is diagonalizable, and $Q\Lambda Q^T$ is possible because it is symmetric.

Problem 24.3: (8.3 #11.) Complete A to a Markov matrix and find the steady state eigenvector. When A is a symmetric Markov matrix, why is $\mathbf{x}_1 = (1, \dots, 1)$ its steady state?

$$A = \begin{bmatrix} .7 & .1 & .2 \\ .1 & .6 & .3 \\ _ & _ & _ \end{bmatrix}.$$

Solution: Matrix A becomes:

$$A = \begin{bmatrix} .7 & .1 & .2 \\ .1 & .6 & .3 \\ .2 & .3 & .5 \end{bmatrix},$$

with steady state vector $(1,1,1)$. When A is a *symmetric* Markov matrix, the elements of each row sum to one. The elements of each row of $A - I$ then sum to zero. Since the steady state vector \mathbf{x} is the eigenvector associated with eigenvalue $\lambda = 1$, we solve $(A - \lambda I)\mathbf{x} = (A - I)\mathbf{x} = \mathbf{0}$ to get $\mathbf{x} = (1, \dots, 1)$.