

18.06 Session 4

Problem 4.1: What matrix E puts A into triangular form $EA = U$? Multiply by $E^{-1} = L$ to factor A into LU .

$$A = \begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

Solution: We will perform a series of row operations to transform the matrix A into an upper triangular matrix. First, we multiply the first row by 2 and then subtract it from the second row in order to make the first element of the second row 0:

$$\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 2 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 3 & 0 \\ 0 & -2 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

Next, we multiply the first row by 2 (again) and subtract it from the third row in order to make the first element of the third row 0:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 \\ 0 & -2 & 0 \\ 2 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 3 & 0 \\ 0 & -2 & 0 \\ 0 & -6 & 1 \end{bmatrix}$$

Now, we multiply the second row by 3 and subtract it from the third row in order to make the second element of the third row 0:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 \\ 0 & -2 & 0 \\ 0 & -6 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 3 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{bmatrix} = U.$$

We take the three matrices we used to perform each operation and multiply them to get E :

$$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 4 & -3 & 1 \end{bmatrix} = E.$$

To check, we evaluate EA :

$$\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 4 & -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 2 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 3 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{bmatrix} = U.$$

To find E^{-1} , use the Gauss-Jordan elimination method:

$$\begin{aligned} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ -2 & 1 & 0 & 0 & 1 & 0 \\ 4 & -3 & 1 & 0 & 0 & 1 \end{array} \right] &\longrightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 2 & 1 & 0 \\ 0 & -3 & 1 & -4 & 0 & 1 \end{array} \right] \\ &\longrightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 2 & 1 & 0 \\ 0 & 0 & 1 & 2 & 3 & 1 \end{array} \right] \longrightarrow \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 2 & 3 & 1 \end{bmatrix} = E^{-1} \end{aligned}$$

We can check that this is in fact the inverse of E :

$$EE^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 4 & -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 2 & 3 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I.$$

Finally, to factorize A into LU (where $L = E^{-1}$):

$$\begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 2 & 0 & 1 \end{bmatrix} = A = LU = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 2 & 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Problem 4.2: (2.6 #13. *Introduction to Linear Algebra: Strang*) Compute L and U for the symmetric matrix

$$\mathbf{A} = \begin{bmatrix} a & a & a & a \\ a & b & b & b \\ a & b & c & c \\ a & b & c & d \end{bmatrix}.$$

Find four conditions on a, b, c, d to get $A = LU$ with four pivots.

Solution: Elimination subtracts row 1 from rows 2-4, then row 2 from rows 3-4, and finally row 3 from row 4; the result is U . All the multipliers ℓ_{ij} are equal to 1; so L is the lower triangular matrix with 1's on the diagonal and below it.

$$\begin{aligned} \mathbf{A} &\longrightarrow \begin{bmatrix} a & a & a & a \\ 0 & b-a & b-a & b-a \\ 0 & b-a & c-a & c-a \\ 0 & b-a & c-a & d-a \end{bmatrix} \longrightarrow \begin{bmatrix} a & a & a & a \\ 0 & b-a & b-a & b-a \\ 0 & 0 & c-b & c-b \\ 0 & 0 & c-b & d-b \end{bmatrix} \\ &\longrightarrow \begin{bmatrix} a & a & a & a \\ 0 & b-a & b-a & b-a \\ 0 & 0 & c-b & c-b \\ 0 & 0 & 0 & d-c \end{bmatrix} = U, L = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix}. \end{aligned}$$

The pivots are the nonzero entries on the diagonal of U . So there are four pivots when these four conditions are satisfied: $a \neq 0, b \neq a, c \neq b$, and $d \neq c$.