

$$F = C_1rv + C_2r^2v^2 \quad F = \frac{mMG}{r^2} \quad \mathbf{F} = \frac{d\mathbf{p}}{dt} = m\mathbf{a} \quad dW = \mathbf{F} \cdot d\mathbf{r} \quad a_{cent} = \frac{v^2}{r} = \omega^2r$$

$$U = \frac{-mMG}{r} \quad U = mgh \quad U = \frac{1}{2}kx^2 \quad K = \frac{1}{2}mv^2 \quad K = \frac{1}{2}I\omega^2$$

$$E_{tot} = K + U = \frac{1}{2}mv^2 - \frac{mMG}{r} = \frac{-mMG}{2a} \quad \mathbf{L} = \mathbf{r} \times \mathbf{p} \quad I = \sum_i m_i r_i^2$$

$$m_1r_1 = m_2r_2 \quad v = \omega r \quad T^2 = \frac{4\pi^2(r_1 + r_2)^3}{G(m_1 + m_2)} \quad \omega = \frac{d\theta}{dt} \quad \omega = \sqrt{k/m}$$

$$\boldsymbol{\tau} = \mathbf{r} \times \mathbf{F} = I\boldsymbol{\alpha} = \frac{d\mathbf{L}}{dt} \quad \alpha = \frac{d\omega}{dt} \quad T = \frac{2\pi}{\omega} \quad L = I\omega \quad \mathbf{I} = \int_0^{\Delta t} \mathbf{F} dt = \mathbf{p}_f - \mathbf{p}_i$$

$$\omega = \sqrt{g/l} \quad \omega_{pr} = \frac{\tau}{L_s} \quad T^2 = \frac{4\pi^2a^3}{GM}$$


Solid disk of mass M and radius R rotating about its cylindrical axis: $I = \frac{1}{2}MR^2$

$$v_f - v_i = -u \ln\left(\frac{m_f}{m_i}\right) - gt \quad I = I_{cm} + Md^2 \quad I_z = I_x + I_y$$

$$f' = f \left(1 + \frac{v}{c} \cos \theta\right) \quad \lambda' = \lambda \left(1 - \frac{v}{c} \cos \theta\right)$$

Problem 1 (35 points)


An unknown mass, m_1 , hangs from a massless string and descends with an acceleration $g/2$. The other end is attached to a mass m_2 which slides on a frictionless horizontal table. The string goes over a uniform cylinder of mass $m_2/2$ and radius R (see figure). The cylinder rotates about a horizontal axis without friction and the string does not slip on the cylinder. Express your answers in parts b, c, and d in terms of g , m_2 , and R .



- (8) Draw free-body diagrams for the cylinder and the two masses.
- (9) What is the tension in the horizontal section of the string?
- (9) What is the tension in the vertical section of the string?
- (9) What is the value of the unknown mass m_1 ?

Problem 2 (30 points)

A bullet of mass m_1 is fired into a pendulum of mass m_2 and length L . The speed of the bullet as it enters the mass m_2 is V_1 (see figure).



First, assume that the collision is elastic, and that $m_1 \ll m_2$.


- (6) If the pendulum is initially at rest, what is the speed of the bullet after the collision?
- (8) Now suppose that when the collision occurs, the pendulum, at the bottom of its swing, is moving to the left with velocity V_2 . What now is the speed of the bullet after the elastic collision?

Now assume that the collision is completely inelastic. The pendulum is at rest before the collision, $m_1 < m_2$, but the speed V_1 of the bullet is unknown.

- (8) After the collision the pendulum moves to the right and it comes to a halt when the string makes an angle θ_{\max} with the vertical. What was the speed of the bullet? Substitute in your answer $\theta_{\max} = 0$. Does your result make sense?
- (8) Could θ_{\max} be 90° ? Explain your answer.

Problem 3 (35 points)

A solid, uniform disk of mass M and radius R is oscillating about an axis through P . The axis is perpendicular to the plane of the disk. Friction at P is negligibly small and can be ignored. The distance from P to the center, C , of the disk is b (see figure). The gravitational acceleration is g .



- (7) When the displacement angle is θ , what then is the torque relative to point P ?
- (7) What is the moment of inertia for rotation about the axis through P ?
- (7) The torque causes an angular acceleration about the axis through P . Write down the equation of motion in terms of the angle θ and the angular acceleration.

As the disk oscillates, the maximum displacement angle, θ_{\max} , is very small, and the motion is a near perfect simple harmonic oscillation.

- (7) What is the period of oscillation?
- (7) As the disk oscillates, is there any force that the axis at P exerts on the disk? Explain your answer.