# Lecture 1: The geometry of linear equations

The fundamental problem of linear algebra is to solve n linear equations in n unknowns; for example:

$$2x - y = 0 \\
-x + 2y = 3.$$

In this first lecture on linear algebra we view this problem in three ways.

The system above is two dimensional (n = 2). By adding a third variable z we could expand it to three dimensions.

### **Row Picture**

Plot the points that satisfy each equation. The intersection of the plots (if they do intersect) represents the solution to the system of equations. Looking at Figure 1 we see that the solution to this system of equations is x = 1, y = 2.




Figure 1: The lines 2x - y = 0 and -x + 2y = 3 intersect at the point (1,2).

We plug this solution in to the original system of equations to check our work:

$$\begin{array}{rcl} 2 \cdot 1 - 2 & = & 0 \\ -1 + 2 \cdot 2 & = & 3. \end{array}$$

The solution to a three dimensional system of equations is the common point of intersection of three planes (if there is one).

#### Column Picture

In the column picture we rewrite the system of linear equations as a single equation by turning the coefficients in the columns of the system into vectors:

$$x \begin{bmatrix} 2 \\ -1 \end{bmatrix} + y \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \end{bmatrix}.$$

Given two vectors  $\mathbf{c}$  and  $\mathbf{d}$  and scalars x and y, the sum  $x\mathbf{c} + y\mathbf{d}$  is called a *linear combination* of  $\mathbf{c}$  and  $\mathbf{d}$ . Linear combinations are important throughout this course.

Geometrically, we want to find numbers x and y so that x copies of vector  $\begin{bmatrix} 2 \\ -1 \end{bmatrix}$  added to y copies of vector  $\begin{bmatrix} -1 \\ 2 \end{bmatrix}$  equals the vector  $\begin{bmatrix} 0 \\ 3 \end{bmatrix}$ . As we see from Figure 2, x = 1 and y = 2, agreeing with the row picture in Figure 2.




Figure 2: A linear combination of the column vectors equals the vector **b**.

In three dimensions, the column picture requires us to find a linear combination of three 3-dimensional vectors that equals the vector **b**.

#### **Matrix Picture**

We write the system of equations

$$2x - y = 0$$
$$-x + 2y = 3$$

as a single equation by using matrices and vectors:

$$\left[\begin{array}{cc} 2 & -1 \\ -1 & 2 \end{array}\right] \left[\begin{array}{c} x \\ y \end{array}\right] = \left[\begin{array}{c} 0 \\ 3 \end{array}\right].$$

The matrix  $A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$  is called the *coefficient matrix*. The vector  $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$  is the vector of unknowns. The values on the right hand side of the equations form the vector  $\mathbf{b}$ :

$$A\mathbf{x} = \mathbf{b}$$
.

The three dimensional matrix picture is very like the two dimensional one, except that the vectors and matrices increase in size.

#### **Matrix Multiplication**

How do we multiply a matrix A by a vector  $\mathbf{x}$ ?

$$\left[\begin{array}{cc} 2 & 5 \\ 1 & 3 \end{array}\right] \left[\begin{array}{c} 1 \\ 2 \end{array}\right] = ?$$

One method is to think of the entries of x as the coefficients of a linear combination of the column vectors of the matrix:

$$\left[\begin{array}{cc} 2 & 5 \\ 1 & 3 \end{array}\right] \left[\begin{array}{c} 1 \\ 2 \end{array}\right] = 1 \left[\begin{array}{c} 2 \\ 1 \end{array}\right] + 2 \left[\begin{array}{c} 5 \\ 3 \end{array}\right] = \left[\begin{array}{c} 12 \\ 7 \end{array}\right].$$

This technique shows that Ax is a linear combination of the columns of A.

You may also calculate the product Ax by taking the dot product of each row of A with the vector x:

$$\left[\begin{array}{cc} 2 & 5 \\ 1 & 3 \end{array}\right] \left[\begin{array}{c} 1 \\ 2 \end{array}\right] = \left[\begin{array}{c} 2 \cdot 1 + 5 \cdot 2 \\ 1 \cdot 1 + 3 \cdot 2 \end{array}\right] = \left[\begin{array}{c} 12 \\ 7 \end{array}\right].$$

## Linear Independence

In the column and matrix pictures, the right hand side of the equation is a vector **b**. Given a matrix *A*, can we solve:

$$A\mathbf{x} = \mathbf{b}$$

for every possible vector **b**? In other words, do the linear combinations of the column vectors fill the *xy*-plane (or space, in the three dimensional case)?

If the answer is "no", we say that *A* is a *singular matrix*. In this singular case its column vectors are *linearly dependent*; all linear combinations of those vectors lie on a point or line (in two dimensions) or on a point, line or plane (in three dimensions). The combinations don't fill the whole space.